

FIG.1

REPLACEMENT SHEET

2/29




FIG. 2a




FIG. 2b

REPLACEMENT SHEET

3/29

FIG.3a




FIG.3b





FIG.3c



REPLACEMENT SHEET

4/29

FIG.4a




FIG.4b





FIG.4c



5/29

FIG.5 The Distribution of Sex and Age of 100 Patients with A.L.F.

AGE	F	M	TOTAL
1-	2	3	5
11-	1		1
21-	3	5	8
31-	10	2	12
41-	16	7	23
51-	21	7	23
61-	20	1	21
71-	2		2
<u>TOTAL</u>	<u>75</u>	<u>25</u>	<u>100</u>

6/29

ALF Dosage

FIG.6

	Patient		Dosage (cc/q4d) (minimum dose in cc that triggers blastogenesis)	
	No.	%	Range	Mean
Normal	47	47	0.1	0.1
Build up	37	37	0.15-1.0	0.3
Dilution	16	16	0.01- 0.00001	0.05

7/29

FIG.7 The Distribution of Time of Therapy

Months	2	3	4	5	6	7	8	9	10	11	>12
No. Patients	8	14	11	7	7	6	10	6	4	7	20

Average therapy time: 7.4 ± 3.8 months

Range: 2-16 months

8/29

FIG.8 The Distribution of Dosage of Therapy

Dosage (cc)	2	3	4	5	6	7	8	9	10	11	>12
No. Patients	8	13	15	6	8	11	10	6	4	7	14

Average therapy 6.44 ± 3.79 cc

Range: 2-25 cc

9/29

FIG.9 Scale of Clinical Symptoms and Signs of
The Patient After ALF Treatment
(Severity)

Symptoms	Improvement		No Change		Worsening		Total Patients
	No.	%	No.	%	No.	%	
Hypersensitive Reaction	63	63	33	33	1	1	100
Recurrent Infection	38	57	29	43	0	0	67
Fatigue	60	68	27	31	1	1	88
Lack of Concentration	43	54	36	46	0	0	79
Arthritis	19	44	23	54	1	2	43
G.I. Upset	29	40	43	60	0	0	72
Headache	28	44	33	53	2	3	64
Depression	42	58	30	42	0	0	72

10/29

FIG.10 Scale of Clinical Symptoms and Signs of
The Patient After ALF Treatment
(Frequency)

Symptoms	Improvement		No Change		Worsening		Total Patients
	No.	%	No.	%	No.	%	
Hypersensitive Reaction	38	38	62	62	0	0	100
Recurrent Infection	26	39	41	61	0	0	67
Fatigue	38	43	50	57	0	0	88
Lack of Concentration	20	25	59	75	0	0	79
Arthritis	11	26	32	74	0	0	43
G.I. Upset	21	29	51	69	0	0	72
Headache	19	30	45	70	0	0	64
Depression	30	42	42	58	0	0	72

11/29

The Changes of T&B Profile of 92 Patients Treated with ALF

	TOTAL LYMPHOCYTES	T _H	T _H	T _H	T _H /T _B	B _L
BEFORE	2112±632	1624±457	930±45	439±58	2.3±0.8	188±102
AFTER	2232±678 *	1634±544	1027±297 **	478±189 **	1.3±.4	171±120

n = 92

* p<0.05

** p<0.01

12/29

FIG.12 T Lymphocyte Subsets of 92 Patients After AIF Treatment

	TOTAL LYMPHOCYTES		T _H		T _H		T _H	
	NO.	%	NO.	%	NO.	%	NO.	%
INCREASE	52	57	41	46	53	58	55	60
DECREASE	40	43	51	54	39	42	37	40
P	<0.05		>0.05		<0.05		<0.01	

13/29

FIG. 13
CMI Results with ALF Effect
CMI Scores (Number and Size, $\bar{x} \pm SD$)

NO. PATIENTS	BEFORE ALF	AFTER ALF	P
78	5.46 \pm 5.81	9.28 \pm 7.25	<0.001

CMI Positive Score
(Number and Size) After ALF

	NO. PATIENTS	%	P
INCREASE	58	74	<0.001
NO CHANGE	9	12	
DECREASE	11	14	

The Relationship Between CMI and Clinical Symptom and Sign Improvement

	IMPROVEMENT NO.	%	NO IMPROVEMENT NO.	%	P
CMI INCREASE	56	72	2	3	<0.001
CMI NO CHANGE & DECREASE	14	18	6	8	

FIG.14 Side Effects of ALF - (5% of Patients Can Not Tolerate ALF)

1. PAIN AND IRRITATION IN THROAT, BURNING IN THE EYES
2. NAUSEA
3. CHEST TIGHTNESS AND HEART PALPITATION
4. FLU-LIKE SYMPTOMS, HEADACHE, FATIGUE, CHILL
5. INCREASE IN FOOD AND ODOR SENSITIVITY

FIG.15 190 Patients with ALF

	No.	%
Female	131	69
Male	59	31
Age:	2-76 years old	
Mean:	46.8 years old	

FIG. 16

ALF Dosage

Patient		Dosage (cc/q4d) (minimum dose in cc that triggers blastogenesis)	
No.	%	Range	
Normal	88	46	0.1 (1:10)
Build up	84	44	0.15-1.0 (1:10)
Dilution	18	9	0.1 (1:20, 1:10 ² -1:10 ⁶)

FIG.17

The distribution of period of ALF therapy

Month	No. Patients	%
1-3	27	14.2
4-6	19	10.0
7-9	25	13.2
10-12	20	10.5
13-15	11	5.8
16-18	21	11.1
19-21	21	11.1
22-24	7	3.7
24-26	14	7.4
27-29	11	5.8
30-32	5	2.6
33-36	9	4.7

Average: 14.6 months (Range 1-37 months)

FIG. 18

The Distribution of dosage of ALF

Dosage (cc)	No. Patient	%
1-5	45	23.7
6-10	59	31.1
11-20	66	34.7
21-30	19	10
31-40	1	0.5
Total	190	100

Average: 11.27 (cc)

Range: 1-38 (cc)

FIG.19
Scale of Clinical Symptoms and Signs of The C.S. Patient After
ALF Treatment (Severity)

Symptoms	Improvement No. %	No Changes No. %	Worsening No. %	Total Patients
Hypersensitive				
Reaction	51 56.7	33 36.7	6 6.7	90
Recurrent				
Infection	33 50.8	27 41.5	5 7.7	65
Fatigue	58 71.6	17 20.99	6 7.4	81
Lack of				
Concentration	42 57.5	29 39.7	2 2.7	73
Arthritis	20 44.4	22 48.9	3 6.7	45
G.I.Upset	32 44.4	33 45.8	7 9.7	72
Headache	30 40.5	38 51.4	6 8.1	74
Depression	44 63.8	22 31.9	3 4.3	69

FIG.20

Scale of Clinical Symptoms and Signs of The Patient After
ALF Treatment (frequency)

Symptoms	Improvement No. %	No Changes No. %	Worsening No. %	Total Patients
Hypersensitive Reaction	28 36.4	46 59.7	3 3.9	77
Recurrent Infection	27 46.6	27 46.6	4 6.9	58
Fatigue	36 47.4	37 48.7	3 3.9	76
Lack of Concentration	23 33.8	43 63.2	2 2.9	68
Arthritis	13 32.5	24 60.0	3 7.5	40
G.I.Upset	23 37.1	35 56.5	4 6.5	62
Headache	24 38.1	34 54.0	5 7.9	63
Depression	34 54.8	26 41.9	2 3.2	62

FIG.21

T & B Lymphocytes Parameters Changes After ALF

	(N)	Before (mean±SE)	After
WBC	25	4180±404.1**	4828±1164.6
Lymphis	34	1350 ±225.1**	1573.2±448.2
T11	32	989.4±198.1**	1182.6±331.2
T4	30	518.2±107.2**	689.5 ±200.1
Ratio of T4/T8	46	3.62± 0.99**	2.71± 1.04
T8	52	248.6±63.3**	354.3 ±124.9
B4	35	54.8±17.0**	121.3±78.7

**--P<0.01 (Change compared with before ALF therapy)

FIG.22
T&B Lymphocytes parameters improvement after
ALF

	Patients No.	Improved No.	(%)
WBC	25	18	72%
Lymphs	34	25	74%
T11	32	21	66%
T4	30	24	80%
Ratio of T4/T8	46	37	80.4%
T8	52	40	76.9%
B4	35	27	77%

FIG.23

CMI Improvements of 93 Patients with ALF

<u>Total No. of Antigen Responses</u>	
Increase	247*
Decrease	171
<u>No change</u>	<u>233</u>
Total	651

* P<0.001 compared with decrease

FIG.24**Diagnosis**

	<u>No.</u>	<u>%</u>
Chemical sensitivity	127	67
Food Sensitivity	106	56
Inhalant Sensitivity	84	44
EMF Sensitivity	9	5

25/29

FIG.25

Diagnosis

	No.	%
N.S	104	54.7
C.V	53	27.9
G.I	59	31.1
G.U	15	7.9
M.S	107	56.3
EYE	4	2.1
Respiratory	87	45.8
Skin	22	11.6
Endocrine	24	12.6
Immune		
dysregulation	99	52.1
Chronic		
infectioin	42	22.1
Implant	7	3.7
Parasite	3	1.6
Carcinoma	3	1.6
Candidosis	8	4.2
HIV positive	1	0.5

FIG.26

50 Year Old White Female
HIV Positive

0.1cc every 4 doses

CMI before -1	CMI after 3
WBC - 3,500	10,000
T _H 964	1359
T _H * 54	73%
T ₄ 149	750
T ₄ * 8%	40%
T ₈ 464	875
T ₈ * 26	47%
T ₄ /T ₈ .17	1.62
B 36	93
B% 2	5%

Sx-fatigue,
neuropathy
Stomach Pain
Non healing toe
Lesion for 2 yrs.

Energy
- gone
- gone
- gone
- gone

Took ALF for 5 months, then stopped
All Sx exacerbated for 3 weeks.
Restarted ALF and Sx disappeared

REPLACEMENT SHEET

26/29

REPLACEMENT SHEET

27/29

FIG.27

B: Before ALF

A: After ALF

PATIENTS WITH CANCER & ALF

	46 Y F Breast Cancer	59 Y F Breast Cancer	84 Y F Leukemia	67 Y F para- thyroid tumor	42 Y F Breast Cancer	63 Y F Left Breast Mastectomy	23 Y M Osteo- genic Sarcoma
	3/1/95 9/3/96	6/1/95 7/21/95	10/20/94 2/7/95	3/25/96 1/9/97	10/10/94 5/30/95	10/31/96 1/10/97	4/8/96 9/24/96
	0.1cc 0.2cc 90 doses	0.1cc 40	0.1cc 60	1:10 ³ -1:10 0.1cc 130	0.1cc 90	0.1cc 70	0.1cc 50
Tetanus	B 3	B 0	B A	B A	B A	B A	B A
Diphtheria	4	0	6	4	0	0	0
Streptococcus	0	0	0	0	0	0	0
Tuberculin	0	0	0	0	0	0	0
Control	0	0	0	0	0	0	0
Candida	0	0	0	0	0	0	0
Trichophyton	0	0	0	0	0	0	0
Proteus	0	4	4	0	4	0	0

REPLACEMENT SHEET

28/29

FIG.28

	46 Y F Breast Cancer	59 Y F Breast Cancer	84 Y F Leuk- emia	67 Y F Para- thyroid tumor	42 Y F Breast Cancer	61 Y F Left Breast	23 Y M Osteo- sarcoma
NBC	B A 6900 4300↓ 5100 4200↓	B A 22300↑	B A 7000	B A 4700 4500 6700	B A 5300	B A 6700	B A
Lymphocytes	2277 1720 1848	15387↑	2590	1833 1305 2412			
Ly†	33 50 44	69	37	27	39 29 36		
T ₁	1867 1056↓ 1734 1478	4923↑	1868	1188 861 1787			
T ₁ †	82 62 80	32	76	83	79 66 72		
T ₁	1002 3784 832	2308↑	1606	587	733 469 724		
T ₁ †	44 221 45	154	62↑	41	40 36 30		
T ₁	433 482 591	2616↑	544	329	495 235 699		
T ₁ †	19 28 32	17	21	23	27 18 29		

REPLACEMENT SHEET

29/29

FIG.29

T ₁ /T ₂	15	2.32 0.794 1.41	0.94	2.951	1.5	2	1.03
S ₁	332	205 120 92	18481	104	155	184	241
S ₂	13	8 7 5	12	4	9	8	10
S: Severity							
F: Frequency							
Allergic	S	A	A	A	0	0	0
	F	↑	↑	↑	0	0	0
Headache	S	0	0	0	0	0	0
	F	0	0	0	0	0	0
Recurrent	S	↑	↑	↑	0	0	0
	F	↑	↑	↑	0	0	0
Fatigue	S	↑	↑	↑	0	0	0
	F	↑	↑	↑	0	0	0
Difficulty	S	↑	↑	↑	0	0	0
	F	0	0	0	0	0	0
Concen- tration	S	↑	↑	↑	0	0	0
	F	↑	↑	↑	0	0	0
Arthritis	S	↑	↑	↑	0	0	0
	F	↑	↑	↑	0	0	0
Of Upset	S	↑	↑	↑	0	0	0
	F	↑	↑	↑	0	0	0
Depression	S	↑	↑	↑	0	0	0
	F	↑	↑	↑	0	0	0